
No.



9100202

# THE CONTRES OF THE CONTRES OF THE CONTRESS OF

# TO ALL TO WHOM THESE PRESENTS SHALL COME: University of Illinois

Withereas, there has been presented to the

#### Secretary of Agriculture

AN APPLICATION REQUESTING A CERTIFICATE OF PROTECTION FOR AN ALLEGED NOVEL VARIETY OF SEXUALLY REPRODUCED PLANT, THE NAME AND DESCRIPTION OF WHICH ARE CONTAINED IN THE APPLICATION AND EXHIBITS, A COPY OF WHICH IS HEREUNTO ANNEXED AND MADE A PART HEREOF, AND THE VARIOUS REQUIREMENTS OF LAW IN SUCH CASES MADE AND PROVIDED HAVE BEEN COMPLIED WITH, AND THE TITLE THERETO IS, FROM THE RECORDS OF THE PLANT VARIETY PROTECTION OFFICE, IN THE APPLICANT(S) INDICATED IN THE SAID COPY, AND WHEREAS, UPON DUE EXAMINATION MADE, THE SAID APPLICANT(S) IS (ARE) ADJUDGED TO BE ENTITLED TO A CERTIFICATE OF PLANT VARIETY PROTECTION UNDER THE LAW.

NOW, THEREFORE, THIS CERTIFICATE OF PLANT VARIETY PROTECTION IS TO GRANT UNTO THE SAID APPLICANT(S) AND THE SUCCESSORS, HEIRS OR ASSIGNS OF THE SAID APPLICANT(S) FOR THE TERM OF eighteen years from the date of this grant, subject to the payment of the required fees and periodic replenishment of viable basic seed of the variety in a public repository as provided by LAW, the right to exclude others from selling the variety, or offering it for sale, or reproducing it, importing it, or exporting it, or using it in producing a hybrid or different

TY THEREFROM, TO THE EXTENT PROVIDED BY THE PLANT VARIETY PROTECTION ACT,
UNITED STATES SEED OF THIS VARIETY (1) SHALL BE SOLD BY VARIETY NAME ONLY AS
OF CERTIFIED SEED AND (2) SHALL CONFORM TO THE NUMBER OF GENERATIONS
THE OWNER OF THE RIGHTS. (84 STAT. 1542, AS AMENDED, 7 U.S.C. 2321 ET SEQ.)

WHEAT

'Howell'

In Esstimony Winercot, I have hereunto set my hand and caused the seal of the Plant Variety Protection Office to be affixed at the City of Washington, D.C.

this 30th day of April in the year of our Lord one thousand nine bundred and ninety-three.

Allest

Kennett Herans

Commissioner

Plant Variety Protection Office Agricultural Marketing Service City Estary of Agriculture

Public reporting burden for this collection of information is estimated to average 30 minutes per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Department of Agriculture, Clearance Office, OIRM, Room 404-W, Washington, D.C. 20250; and to the Office of Management and Budget, Paperwork Reduction Project (OMB #0581-0055), Washington, 20250.

FORM APPROVED: OMB 0581-0055, Expires 1/31/91

U.S. DEPARTMENT OF AGRICULTURAL MARK	AGRICULTURE		Application is required in order to
APPLICATION FOR PLANT VARIE	TY PROTECTION	CERTIFICATE	determine if a plant variety protection certificate is to be issued (7 U.S.C. 2421) Information is held confidential unti- certificate is issued (7 U.S.C. 2426).
NAME OF APPLICANT(S) (as it is to appear on the Certificate)		2. TEMPORARY DESIGNATION OR EXPERIMENTAL NO.	3. VARIETY NAME
University of Illinois		IL 82-3298	Howell
4. ADDRESS (street and no. or R.F.D. no., city, state, and ZIP)	<u> </u>	5. PHONE (Include area code)	FOR OFFICIAL USE ONLY
Illinois Agricultural Experiment St 1301 W. Gregory 211 Mumford Hall University of Illinois Urbana, IL 61801	ation	(217) 333-0240	9100202 F Date
6. GENUS AND SPECIES NAME	7. FAMILY NAME (Botanio	ai)	1 June 20, 1991
Triticum aestivum L.	Graminae		G A.M. P.M.
8. CROP KIND NAME (Common Name)	<u> </u>	PATE OF DETERMINATION	F Filing and Examination Fee:
Cofe Dod History III.		and the second second	E \$ 2150.°°
Soft Red Winter Wheat  10. IF THE APPLICANT NAMED IS NOT A "PERSON," GIVE FORM OF ORGA	ANIZATION (Correction and	8/88	S Date
University of Illinois	NIZATION (Corporation, part	ersnip. association, etc.)	R Sure 20,1991 Certificate Fee:
11. IF INCORPORATED, GIVE STATE OF INCORPORATION	12. DA	TE OF INCORPORATION	f \$250.00
			\$ Opr.9, 1993
University of Illinois 211 Mumford Hall, 1301 West Gregory  14. CHECK APPROPRIATE BOX FOR EACH ATTACHMENT SUBMITTED (Fo.  a. X Exhibit A, Origin and Breeding History of the Variety.  b. X Exhibit B, Novelty Statement.  c. X Exhibit C, Objective Description of Variety.  d. X Exhibit D, Additional Description of Variety.  e. X Exhibit E, Statement of the Basis of Applicant's Ownersh 1. X Seed Sample (2,500 viable untreated seeds). Date Seed 9. X Filing and Examination Fee (\$2,150) made payable to "  15. DOES THE APPLICANT(S) SPECIFY THAT SEED OF THIS VARIETY BE SO Protection Act.)  X YES (If "YES," answer items 16 and 17 be NUMBER OF GENERATIONS?	nip. I Sample mailed to Plant V. Treasurer of the United Sta DLD BY VARIETY NAME ONLY AND NO (# "NO. TO 17. IF "YES" TO	ariety Protection Office 5/29/9 tes." AS A CLASS OF CERTIFIED SEED? (See., "skip to item 18 below) ITEM 16, WHICH CLASSES OF PRODU	2 section 83(a) of the Plant Variety  ICTION BEYOND BREEDER SEED?
X YES  NO	1	DATION X REGIST	ERED X CERTIFIED
18. DID THE APPLICANT(S) PREVIOUSLY FILE FOR PROTECTION OF THE VALUE O	Patent Act. Give date	-	
20. The applicant(s) declare(s) that a viable sample of basic se request in accordance with such regulations as may be appl. The undersigned applicant(s) is (are) the owner(s) of this uniform, and stable as required in section 41, and is entitle Applicant(s) is (are) informed that false representation here.	icable. sexually reproduced no d to protection under the	ovel plant variety, and believe provisions of section 42 of the F	o(s) that the variety is distinct
SIGNATURE OF APPLICANT [Owner(s)]	DATE		
1 4 Dock	Men	elor	900/9/
SIGNATURE OF APPLICANT (Gwner(s))	CAPACITY OR TIT	LE	DATE

FORM CSSD-470 (5-89) Edition of FORM LS-470, 3-86, is obsolete.

#### Exhibit A

#### Origin and Breeding History of Howell

- 1) Howell (previously designated IL 82-3298) was developed by the Illinois Agricultural Experiment Station. The parents of Howell are: McNair 48-23 / IL 70-2225 // CI 13855 /3/ Arthur / Blueboy // TN 1571. Howell was first selected in 1982 as a headrow from a single F<sub>4</sub> plant. Howell was repurified in 1986 and 1988. Heads selected from Howell were grown in headrows in 1986, and headrows with variant types in them were removed. The remaining headrows were harvested in bulk. Heads were again selected in 1987, and 200 headrows were grown in 1988. Most of the rows were uniform and similar to each other. Any rows that differed were removed. Seed from the remaining rows was bulked.
- 2) Breeder seed of Howell was produced from the progeny of the bulked headrows grown in 1988. The seed was increased in 1989. The field was inspected frequently and variants rogued. This breeder seed  $(F_{13})$  was increased in 1990 and variants again rogued. Foundation seed is in production in 1991 from this repurified breeder seed.
- 3) Howell has been stable and true-breeding; however, up to 0.5 % other types, including tall variants which are 6 to 12 cm taller than most plants in the variety, are allowed. Except for the difference in height, the tall variants are very similar in appearance to other plants in the variety; however, some variants may not be awned.
- 4) The performance of Howell was tested in breeding nursery trials in Illinois from 1985 1990, in variety testing trials in Illinois from 1985 1990, and for three years in the Uniform Eastern Soft Red Winter Wheat Nursery from 1986 1988.
- 5) Howell is high-yielding, has excellent winter hardiness, and has very high test weight. Howell was selected in comparison to popular varieties in Illinois, including Caldwell and Cardinal.

#### Exhibit B

#### **Novelty Statement**

Howell is an awned white-chaffed soft red winter wheat cultivar. It is similar to Cardinal in maturity and height and yields equal to or better than Cardinal in Illinois. It is easily distinguished from Cardinal because Howell is awned and Cardinal is apically awnletted. Howell stands well. Howell has excellent winter hardiness and excellent test weight. Test weight of Howell has averaged 2.5 pounds per bushel more than Cardinal in 18 tests in Illinois.

Howell can be distinguished from many soft red winter wheat varieties because it is awned. The awned variety, Dynasty, can be distinguished from Howell by comparing height and maturity. Dynasty is shorter and earlier than Howell.

Howell is moderately resistant to soil borne wheat mosaic virus, barley yellow dwarf virus, and stem rust (*Puccinia graminis*, Pers. f. sp. *tritici* Eriks. & Henn.) Howell is moderately susceptible to leaf rust (*Puccinia recondita*, Rob. ex Desm. f. sp. *tritici*) and powdery mildew (*Erysiphe graminis*, DC. f. sp. *tritici* E. Marchal). Howell is susceptible to biotypes B and D of the Hessian fly (*Mayetiola destructor*, Say). It has not been evaluated with other biotypes of Hessian fly.

Howell' is most similar to 'Cardinal!.

AAA-(per letter)

10 Mar

University of Il ois at Urbana-Champaign

Department of Agronomy

W-203 Turner Hall 1102 South Goodwin Avenue Urbana, IL 61801-4798

217 333-3420 217 333-9817 fax 9100202

February 25, 1993

Alan A. Atchley Plant Variety Protection Office NAL Bldg., Room 500 10301 Baltimore Blvd. Beltsville, MD 20705-2351

Dear Alan:

As one of the breeders responsible for the development and description of 'Howell' soft red winter wheat, I am responding to your questions to Dr. Holt regarding Howell.

Howell has been stable and true-breeding over all  $(F_5 \text{ to } F_{13})$  generations since the original selection in the  $F_4$ . The original seed increase of Howell contained some tall variants. The repurifications in 1986 and 1988 were carried out to reduce the frequency of these tall variants and make the variety more homogeneous for height. The selections in 1986 and 1988 were made based on plant height. Headrows that were taller or shorter than the majority of the headrows were removed, and seed of the remaining headrows was bulked to provide seed to increase.

In reference to the novelty statement, Howell is most similar to the variety Cardinal. I did not make that clear in the original novelty statement. Howell is distinguished from Cardinal because Cardinal is apically awnletted, but Howell is awned. Also, Howell can also be distinguished from Cardinal on the basis of coleoptile color (Exhibit C, item 13). Cardinal has a purple coleoptile, and Howell has a white coleoptile.

The inconsistency related to the description of Howell's reaction to stem rust, leaf rust and powdery mildew is related to the Exhibit C form because the form provides only 2 classes (susceptible or resistant). I concur with you that the meaning of "susceptible", "moderately susceptible" and "moderately resistant" are not the same. I believe Howell should be classified as moderately resistant to stem rust and moderately susceptible to leaf rust and powdery mildew. I've included that information in the novelty statement, but it is probably more appropriate to have it included in Exhibit D, additional description of the variety. I consider these descriptions in the text to be more accurate than the information from Exhibit C.

nosification Shibat

#### U.S. DEPARTMENT OF AGRICULTURE AGRICULTURAL MARKETING SERVICE LIVESTOCK AND SEED DIVISION BELTSVILLE, MARYLAND 20705

EXHIBIT C

# OBJECTIVE DESCRIPTION OF VARIETY WHEAT (TRITICUM SPP.)

University of Illinois  Address (Street and No. or R.P.D. No., City, State, and EIP Code)  Illinois Agricultural Experiment Station  1301 W. Gregory  1311 Muniford Hall, University of Illinois, Urbana, IL 61801  Place the appropriate number that describes the varietal character of this variety in the boxes below.  Place a zero is first box (e-a- [0] 1 o [0] o [0] ) when number is either 99 or less on 9 or less.  1, KIND.  1   = COMMON 2 = DUNUM 3 = EMMER 4 = SPELT 5 = POLISH 6 = POULARD 7 = CLUB  2, TYPE.  2   I = SPRING 2 = WINTER 3 = OTHER (Speelty)  1   2 = MARD  2   I = SPRING 2 = WINTER 3 = OTHER (Speelty)  1   2 = MARD  2   I = WINTER 2 = RED 3 = OTHER (Speelty)  1   2 = ARTHUR 2 = SCOUT 3 = CHRIS  1   A = LEMH	NAME OF APPLICANTIE	
ADDRESS (Street and Mo. or R.P.D. Mo., City, State, and EPP Code)  1301 W. Gregory  211 Mumford Hall, University of Illinois, Urbana, IL 61801  1301 W. Gregory  212 Mumford Hall, University of Illinois, Urbana, IL 61801  1301 W. Gregory  213 Mumford Hall, University of Illinois, Urbana, IL 61801  1302 W. Gregory  214 Mumford Hall, University of Illinois, Urbana, IL 61801  1303 W. Gregory  215 Mumford Hall, University of Illinois, Urbana, IL 61801  1304 W. Gregory  216 Mumford Hall, University of Illinois, Urbana, IL 61801  1305 W. School W.	·	
Thinois Agricultural Experiment Station  301 W. Gregory  211 Mumford Hall, University of Illinois, Urbana, IL 61801  Place the appropriate number that describes the varietal character of this variety in the boses below.  Place a zero in first box (***-[0] 8   7] or [0] 9) when number is clither 99 or leas or 9 or leas.  1. KING:  1 1 = COMMON 2 = DURRUM 3 = EMMER 4 = SPELT 5 = POLISH 6 = POULARO 7 = CLUB  2 YYPE.  2 1 = SPRING 2 = WINTER 3 = OTHER (Specify)  1 2 = MARD  2 = OTHER (Specify)  3 = OTHER (Specify)  1 2 = MARD  3 = OTHER (Specify)  3 = OTHER (Specify)  1		7100202
1301 W. Gregory 211 Mumford Hall, University of Illinois, Urbana, IL 61801  Place the appropriate number that describes the varietal character of this variety in the boxes below.  Place a zero in first box (e-s-		
211 Mumford Hall, University of Illinois, Urbana, IL 6180  Place the appropriate number that describes the varietal character of this variety in the boxes below.  Place a zero in first box (*-4- [0] 8] 7) or [0] 7) when number is either 99 or leas or 9 or leas.  1. KIND:  1	1301 W. Gregory	DEHGNATION
Place the appropriate number that describes the varietal character of dala variety in the boxes below.  Place a zero is first box (*-4- 0 8 9 9 or 0 9 ) when number is either 99 or less or 9 or less.  1. KIND;  1. I = COMMON 2 = DURUM 3 = EMMER 4 = SPELT 5 = POLISH 6 = POULARD 7 = CLUB  2. TYPE.  2. I = SPRING 2 = WINTER 3 = OTHER (Specify) 1 2 = HARD  1. SPRING 2 = WINTER 3 = OTHER (Specify) 2 = HARD  2. I = WINTER 2 = RED 3 = OTHER (Specify) 1 2 = HARD  3. SEASON - HUMBER OF DAYS FROM EMERGENCE TO:  2. 2 9 FIRST FLOWERING  2. MATURITY (SOS Flowering):  1. MATURITY (SOS FLOWERING):  2. MATURITY (SOS FLOWERING):  3. SEASON - MUMBER OF FLOWERING  3. SEASON - MUMBER OF FLOWERING  3. SEASON		bana, IL 61801
1. KIND: 1 1= COMMON 2 = DURUM 3 = EMMER 4 = SPELT 5 = POLISH 6 = POULARD 7 = CLUB 2. TYPE: 2 1 = SPRING 2 = WINTER 3 = OTHER (Specify) 3. SEASON - NUMBER OF DAYS FROM EMERGENCE TO: 2 2 9 FIRST FLOWERING 2	Place the appropriate number that describes the varietal character	of this variety in the boxes below.
2 TYPE: 2 I = SPRING 2 = WINTER 3 = OTHER (Specify) 1 2 = HARD 2 I = WHITE 2 = RED 3 = OTHER (Specify) 2 I = WHITE 2 = RED 3 = OTHER (Specify) 2 I = WHITE 2 = RED 3 = OTHER (Specify) 3 SEASON - NUMBER OF DAYS FROM EMERGENCE TO: 2 2 9 FIRST FLOWERING  1 MO. OF DAYS FROM EMERGENCE TO: 2 3 4 LAST FLOWERING  1 MO. OF DAYS FROM EMERGENCE TO: 2 3 4 LAST FLOWERING  1 MATURITY (SOX Flowering): 1 NO. OF DAYS LATER THAN		
1   SOPT   3   SOTHER (Specify)   1   2   SOPT   3   SOTHER (Specify)   2   1   SOPT   3   SOTHER (Specify)   2   1   SOPT   3   SOTHER (Specify)   3   SOTHER (Specify)   3   SEASON - NUMBER OF DAYS FROM EMERGENCE TO:   2   2   9   FIRST FLOWERING   2   3   4   LAST FLOWERING     1. MATURITY (SOX Flowering):	1 = COMMON 2 = DURUM 3 = EMMER 4 = SPELT 5 =	POLISH 6 = POULARD 7 = CLUB
1 = SPRING   2 = WINTER   3 = OTHER (Specify)   1   2 = HARD	2 TYPE,	
3. SEASON - NUMBER OF DAYS FROM EMERGENCE TO:  2 2 9 FIRST FLOWERING  2 3 4 LAST FLOWERING  1. MATURITY (SOR Flowering):  NO. OF DAYS EARLIER THAN	2 1 = SPRING 2 = WINTER 3 = OTHER (Specify)	
2 2 9 FIRST FLOWERING  1. MATURITY (50% Flowering):    NO. OF DAYS EARLIER THAN   1 = ARTHUR 2 = SCOUT 3 = CHRIS	2   1 = WHITE 2 = RED 3 = OTHER (Specily)	net transfer
MATURITY (50% Flowering):   NO. OF DAYS EARLIER THAN   1	3, SEASON - NUMBER OF DAYS FROM EMERGENCE TO:	
NO. OF DAYS EARLIER THAN    1 = ARTHUR	2 2 9 FIRST FLOWERING	2 3 4 LAST FLOWERING
A = LEMHI S = NUGAINES 6 = LEEDS  PLANT HEIGHT (From soil level to top of head):  10: 7 CM. HIGH CAMECRY 10 107 based on Compared as varieties mentioned in Each.  10: 7 CM. HIGH CAMECRY 10 107 based on Compared as varieties mentioned in Each.  11 = ARTHUR 2 = SCOUT 3 = CHRIS  12 = ARTHUR 2 = SCOUT 3 = CHRIS  13 = ARTHUR 2 = SCOUT 3 = CHRIS  14 = LEMHI 5 = NUGAINES 6 = LEEDS  PLANT COLOR AT BOOTING (See reverse):  11 = YELLOW GREEN 2 = GREEN 3 = BLUE GREEN 1 1 = YELLOW 2 = PURPLE  STEMS  12 Anthocyanin: 1 = ABSENT 2 = PRESENT 1   Vary bloom: 1 = ABSENT 2 = PRESENT    13 Anthocyanin: 1 = ABSENT 2 = PRESENT    14 NO. OF NODES (Originating from node above ground)    15 Anthocyanin: 1 = ABSENT 2 = PRESENT    16 Anthocyanin: 1 = ABSENT 2 = PRESENT    17 ANTHER COLOR:  18 APPLICATION	(, MATURITY (50% Flowering):	
PLANT HEIGHT (From soil level to top of head):  10:7 cm. High CANCES to 10 to per head):  10:7 cm. High CANCES to 10 to per head):  11	NO. OF DAYS EARLIER THAN	1 = ARTHUR 2 = SCOUT 3 = CHRIS
10: 7 CM. HIGH CANCED to 10 to based on wayardon varieties mentioned in Erch. 1  At A /3 March 5  At A /3 March 6  At LEMNI 5 = NUGAINES 6 = LEEDS  PLANT COLOR AT BOOTING (See reverse):  1		1 4 = LEMHI 5 = NUGAINES 6 = LEEDS
CM. SHORTER THAN	1 0: 7 CM. HIGH COWELLS to 107 based on a	comparisons varieties mentioned in Exch. 1
Tarthur   2 = SCOUT   3 = CHRIS		84A/3Mar/935
CM. SHORTER THAN	0 3 CM. TALLER THAN	1
PLANT COLOR AT BOOTING (See reverse):  7. ANTHER COLOR:  1 = YELLOW GREEN 2 = GREEN 3 = BLUE GREEN  1		TEARTHUR 2 = SCOUT 3 = CHRIS
I = YELLOW GREEN 2 = GREEN 3 = BLUE GREEN  Anthocyanin: I = ABSENT 2 = PRESENT  Hairiness of last internode of rachis: I = ABSENT 2 = PRESENT  AND LEAF BELOW  AURICLES:  Anthocyanin: I = ABSENT 2 = PRESENT  I laternodes: I = HOLLOW 2 = SOLID  CM. INTERNODE LENGTH BETWEEN FLAG LEAF  AND LEAF BELOW  AURICLES:  Anthocyanin: I = ABSENT 2 = PRESENT  LEAF:  Flag leaf at I = ERECT 2 = RECURVED booting stage: 3 = OTHER (Specify):  Hairiness: I = ABSENT 2 = TWISTED  2 = PRESENT  4 = ABSENT 2 = PRESENT  5 = ABSENT 2 = PRESENT  4 = ABSENT 2 = PRESENT  5 = ABSENT 2 = PRESENT  4 = ABSENT 2 = PRESENT  5 = ABSENT 2 = PRESENT  4 = ABSENT 2 = PRESENT  5 = ABSENT 2 = PRESENT  5 = ABSENT 2 = PRESENT  6 = ABSENT 2 = PRESENT  7 = ABSENT 2 = PRESENT  7 = ABSENT 2 = PRESENT  9 = ABSENT 2 = PRESENT  9 = ABSENT 2 = PRESENT  1 = ABSENT 2 = PRESENT  2 = ABSENT 2 = PRESENT  2 = PRESENT  3 = ABSENT 2 = PRESENT  4 = ABSENT 2 = PRESENT  7 = ABSENT 2 = PRESENT  9 = ABSENT 2 = PRESENT  1 = ABSENT 2 = PRESENT  2 = ABSENT 2 = PRESENT  1 = ABSENT 2 = PRESENT  2 = ABSENT 2 = PRESENT  1 = ABSENT 2 = PRESENT  1 = ABSENT 2 = P	CM. SHORTER THAN	4 = LEMHI 5 = HUGAINES 6 = LEEDS
STEM:  Anthocyanin: 1 = ABSENT 2 = PRESENT  Hairiness of last internode of rachis: 1 = ABSENT 2 = PRESENT  AND LEAF BELOW  AURICLES:  Anthocyanin: 1 = ABSENT 2 = PRESENT  Anthocyanin: 1 = ABSENT 2 = PRESENT  The state of first leaf sheath: 1 = ABSENT 2 = PRESENT  Anthocyanin: 1 = ABSENT 2 = PRESENT  The state of first leaf sheath: 1 = ABSENT 2 = PRESENT  The state of first leaf sheath: 1 = ABSENT 2 = PRESENT  AND LEAF:  Anthocyanin: 1 = ABSENT 2 = PRESENT  The state of first leaf sheath: 1 = ABSENT 2 = PRESENT  The state of firs	PLANT COLOR AT BOOTING (See reverse): 7.	ANTHER COLOR:
Anthocyanin: 1 = ABSENT 2 = PRESENT  Hairiness of last internode of rachis: 1 = ABSENT 2 = PRESENT  Anthocyanin: 1 = ABSENT 2 = PRESENT  I laternodes: 1 = HOLLOW 2 = SOLID  O 3 CM. INTERNODE LENGTH BETWEEN FLAG LEAF  AND LEAF BELOW  AURICLES:  Anthocyanin: 1 = ABSENT 2 = PRESENT  I Hairiness: 1 = ABSENT 2 = PRESENT  LEAF:  Flag leaf at 1 = ERECT 2 = RECURVED booting stage: 3 = OTHER (Specify):  Hairs of first leaf sheath: 1 = ABSENT 2 = PRESENT  1 Waxy bloom of flag leaf sheath: 1 = ABSENT 2 = PRESENT	2 I = YELLOW GREEN 2 = GREEN 3 = BLUE GREEN	1   1 = YELLOW 2 * PURPLE
Hairiness of last internode of rachis: 1 = ABSENT 2 = PRESENT    Internodes: 1 = HOLLOW 2 = SOLID   4   HO, OF HODES (Originating from node above ground)   0   3   CM. INTERNODE LENGTH BETWEEN FLAG LEAF AND LEAF BELOW  AURICLES:   Anthocyanin: 1 = ABSENT 2 = PRESENT   1   Hairiness: 1 = ABSENT 2 = PRESENT  LEAF:   Flag leaf at 1 = ERECT 2 = RECURVED   2   Flag leaf: 1 = NOT TWISTED 2 = TWISTED   1   Hairiness: 1 = ABSENT 2 = PRESENT   3 = OTHER (Specify):	STEM:	
Hairiness of last internode of rachis: 1 = ABSENT 2 = PRESENT  1   Internodes: 1 = HOLLOW 2 = SOLID  1   MO. OF HODES (Originating from node above ground)  1   AURICLES:  Authoryanin: 1 = ABSENT 2 = PRESENT  1   Hairiness: 1 = ABSENT 2 = PRESENT  LEAF:  Flag leaf at 1 = ERECT 2 = RECURVED booting stage: 3 = OTHER (Specify):  Hairs of first leaf sheath: 1 = ABSENT 2 = PRESENT  2 = PRESENT  2   Flag leaf: 1 = NOT TWISTED 2 = TWISTED  Hairs of first leaf sheath: 1 = ABSENT 2 = PRESENT  2   PRESENT   Twisted   Twis		
AURICLES:  Anthocyanin: 1 = ABSENT 2 = PRESENT  LEAF:  Flag leaf at 1 = ERECT 2 = RECURVED booting stage: 3 = OTHER (Specify):  Hairs of first leaf sheath: 1 = ABSENT 2 = PRESENT  1 Hairs of first leaf sheath: 1 = ABSENT 2 = PRESENT  2 Flag leaf: 1 = NOT TWISTED 2 = TWISTED  1 Waxy bloom of flag leaf sheath: 1 = ABSENT 2 = PRESENT	Hairiness of last internode of rachis: 1 = ABSENT 2 = PRESENT	
Anthocyanin: 1 = ABSENT 2 = PRESENT  LEAF:  Flag leaf at 1 = ERECT 2 = RECURVED booting stage: 3 = OTHER (Specify):  Hairs of first leaf sheath: 1 = ABSENT 2 = PRESENT  1 Hairs of first leaf sheath: 1 = ABSENT 2 = PRESENT  2 Flag leaf: 1 = NOT TWISTED 2 = TWISTED  1 Waxy bloom of flag leaf sheath: 1 = ABSENT 2 = PRESENT	NO. OF HODES (Originating from node above ground)	
LEAF:    Flag leaf at   = RECUT   2 = RECURVED   2   Flag leaf:   1 = NOT TWISTED   2 = TWISTED     Hairs of first leaf sheath:   = ABSENT   2 = PRESENT   2	AURICLES:	
Flag leaf at 1 = ERECT 2 = RECURVED 2 Flag leaf: 1 = NOT TWISTED 2 = TWISTED  Hairs of first leaf sheath: 1 = ABSENT 2 = PRESENT 1 Waxy bloom of flag leaf sheath: 1 = ABSENT 2 = PRESENT	Anthocyania: 1 = ABSENT 2 = PRESENT	Hairiness:   = ABSENT 2 = PRESENT
booting stage: 3 = OTHER (Specify): 2 Flag leaf: 1 = NOT TWISTED 2 = TWISTED  Hairs of first leaf sheath: 1 = ABSENT 2 = PRESENT  Waxy bloom of flag leaf sheath: 1 = ABSENT 2 = PRESENT	LEAF:	
Twaxy bloom of Hag leaf sheats: 12 Abbent 22 Pricesent	booting grane	Flag leaf: 1 = NOT TWISTED 2 = TWISTED
3 MM, LEAF WIDTH (First load below flag load) 2 2 CM. LEAF LENGTH (First load below flag load):	Hairs of first leaf sheath: 1 = ABSENT 2 = PRESENT	Waxy bloom of flag leaf sheath:   * ABSENT 2 = PRESENT
	3 MM, LEAF WIDTH (First leaf below flat leaf)	CM. LEAF LENGTH (First feel below fied feel):

Angle of the second	$\mathbf{r} = \mathbf{r}$		• •
11. HEAD:			
2 Density: 1 = LAX	2 = DENSE	1 Shape: 1 = TAPE 4 = OTHE	RING 2 = STRAP 3 = CLAVATE (R (Specify)
4 Awnedness: 1 = Av	VALESS 2 = APICALLY AWALETED	3 = AWNLETED 4 = AWN	EO
Color at maturity: 5	= WHITE 2 = YELLOW 3 = PINK	4 = RED	
	= BROWN 6 = BLACK 7 = OTH	IER (Specily):	
8. 6 CM. LENGTH		0 9 MM. WIDTH	en e
12. GLUMES AT MATUR	ITY:		
Length: 1 = SHORT		1 2 1	2 = MEDIUM (CA. 3.5 ma)
3 = LONG	(CA. 9 mm.)	J = WIDE (	CA. 4 mm.)
Shoulder 1 = WANT 2 shape: 4 = SQUA	ING 2 = OBLIQUE 3 = ROUNDED RE 5 = ELEVATED 6 = APICULATE	3 Beak: 1 = OBTUS	E 2 = ACUTE 3 = ACUMINATE
13. COLEOPTILE COLOR	R:	14. SEEDLING ANTHOO	YANIN:
1 1 = WHITE 2 = B	ED 3=PURPLE	1 1 = ABSENT	2 = PRESENT
15. JUVENILE PLANT GE	OWTH HABIT:		
2 1 = PROSTRATE	2 = 5EMI-ERECT 3 = ERE	ст	
16. SEED:			
2 Shape: 1 = OVATE	2 = OVAL 3 = ELLIPTICAL	1 Charles 1 - BOUND	DED 2 = ANGULAR
Z 3112pe. 1 - 0 V X 1 Z	1 - OVAL OF LELIF (ICAL	Cheek, 1 - ROUNT	JED I - ANGUEAR
2 Brush: 1 = SHORT	2 = MEDIUM 3 = LONG	1 Bush 1 a voz s	OLLARED 2 = COLLARED
	1 - MEDIUM 3 - LONG	T plant: 1 = NOT C	OLLARED 2 = COLLARED
Phenol reaction (See instructions):	1 = IVORY 2 = FAWN 3 = LT. BROW	YN :	
(See Markettone).	4 = BROWN 5 = BLACK		•
3 Color: 1 = WHITE	2 = AMBER 3 = RED 4 = PURPLE	5 = OTHER (Specify)	
<del></del>		~ / <del>~</del>	as erroneous, 13 Mar 1995
6. 3 MM. LENGTH	3. 7 MM. WIDTH		SEEDS from database;
17. SEED CREASE:			
1 Width: ] = 60% OR L	ESS OF KERNEL 'WINOKA'	Depth: 1 = 20 % O	R LESS OF KERNEL 'SCOUT'
2 = 80% OR L	ESS OF KERNEL 'CHRIS'		R LESS OF KERNEL 'CHRIS'
3 = NEARLY	AS WIDE AS KERNEL 'LEMHI'	3 = 50 % 0	R LESS OF KERNEL 'LEMHI'
18. DISEASE: (0 = Not-Tes	ted, 1 = Susceptible, 2 = Resistant)	,	
1 STEM RUST (Races)	LEAF RUST	O STRIPE RUST	LOOSE SMUT
		<u>-                                    </u>	
1 POWDERY MILDEW	0 BUNT	OTHER (Specify)	
19. INSECT: (0 = Not Tours	d, 1 = Susceptible, 2 = Resistant)	<del></del>	
0 SAWFLY	? APHID (Bydv.)	O GREEN BUG	O CEREAL LEAF BEETLE
OTHER (Specify)			
OTHER (Specity)	HESSIAN FLY	0 GP 0 A	1 B 0 c
	RACES: (		
			[O],
20. INDICATE WHICH YARII	TY MOST CLOSELY RESEMBLES THAT S	UBMITTED:	
CHARACTER	NAME OF VARIETY	CHARACTER	NAME OF VARIETY
Plant tillering	Cardinal	Seed size	Cardinal
Leaf size	Cardinal	Seed shape	Cardinal
Leaf color	Caldwell	Coleoptile elongation	Caldwell
Leaf carriage	Caldwell	Seedling pigmentation	Caldwell

#### INSTRUCTIONS

GENERAL: The following publications may be used as a reference aid for the standardization of terms and procedures for completing this form:

- (a) L.W. Briggle and L. P. Reitz, 1963, Classification of Triticum Species and Wheat Varieties Grown in the United States, Technical Bulletin 1278, United States Department of Agriculture.
- (b) W.E. Walls, 1965, A Standardized Phenol Method for Testing Wheat Seeds for Varietal Purity, contribution No. 28 to the handbook of seed testing prepared by the Association of Official Seed Analysis. (See attachment.)

#### Exhibit D

#### **Additional Description of the Variety**

Coleoptiles of Howell seedlings are white. Stem anthocyanin is absent in Howell. Auricles and ligules and the last rachis internode are glabrous. The heads are fusiform in shape and classified as mid-dense with about 4.5 cm for 10 rachis internodes. Cheeks of the kernels are rounded and the crease is narrow and shallow. The heads, which measure approximately 9 cm in length and 9 mm in width, are primarily erect at maturity. Howell is awned and white-chaffed.

#### Milling and Baking Quality Evaluation of Howell

(Data from USDA Soft Wheat Quality Laboratory, Wooster, Ohio)

Howell has exhibited good to excellent milling quality (Tables 1-5). For example, flour yield and break flour yield for Howell have been high. The high test weight of Howell also contributes to good milling quality.

Although the baking quality of Howell has not been as good as the milling quality, the baking quality of Howell is equivalent to some varieties that are currently in commercial production (Tables 1-5). For example, the 1989 AWRC for Howell was similar to Tyler and Clark, and the 1989 cookie diameter was similar to Auburn and Clark (Table 5). Howell generally has slightly higher protein percentage than some soft red winter wheat varieties. It has also had slightly lower cookie diameter and cake volume than some varieties. Nevertheless, because of the diverse end uses for soft red winter wheat, the baking quality of Howell should be acceptable to the industry.

Table 1. Wheat milling and baking quality data from samples grown in Illinois in 1985.

SAMPLE NO			MILLI		BAKING QUALITY	COMBINED	ADJ. YIELD	PROTEIN	AHRC	SOFT- NESS		ALLIS E	QUIVALEN	TS .
190			SCOR		SCORE	SCORE	,		******	EQUIV.		ESI	BR FL	EXT.
		STANDOON - Softy							F0 F				27.6	75.5
****		אירועויוני /	100	A	100 A	100 A	72.8	10.9	52.5	48.0	1	12	35.4	76.8
****		BENCHMARK	104.5		106.5 A	104.5 A	74.1	8.3	52.4,	57.7	1	10.2	30.9	75.3
553 <b>5</b>	1	MGH 10950		8	98.9 B	98.9 B	72.6	10.5	53.8	52.1	1	12.3		74.7
5536	2	MOH 10435		B	93.1 C	93.1 C	72.0	1 <b>i</b>	54.6	51.4		13.1	30.3	
5537	3	MOW 11365	98.6	В	101.6 A	98.6 8	72.4	11.1	52.9	55.3	ł	12.6	33.5	75. 1
5538	4	MOW 11165	97	В	86.6 D	86.6 D	71.9*	12.2*	55 t	50.3	1	13.2	29.5	74.6
5539	5	MOW 11081	95.7	B	89.2 D	89.2 D	71.5*	11	55.7 <b>*</b>	53.4	ł	13.8	32	74.2
5540	. 6	IL 81-3210	99. 1	B	98 B	98 B	72.5	11. i	53.6	52.2	ı	12.4	31	75.2
5541	7	IL-81-3696	95.6	9	98.7 B	95.6 B	71.5 <del>*</del>	12.6*	52.5	52.5	ŧ	13.8	31.2	74.2
5542	8	IL 81-3786	98.7	B	95.9 B	95.9 B	72.4	12.2*	53.4	53.4	ł	12.5	31.9	<b>75.</b> i
5543	9	IL 81-2766	103.7	A	91 C	91 C	73.9	11.7	54.2	49.3	i	10.5	28.7	76.5
5544	10	IL 82-3298	96.2		86.1 D	86.1 D	71.7*	11	55.6*	48.6	1_	13.6	28.1	74.4
5545	11	IN 69195C9-4-1-1-38-	97.3		96.3 B	96.3 B	72.0*	11.5	53.5	51.0	1	13.1	30	74.7
5546	12	IN 76148A2-10-4-3	96	B	101.4 A	96 B	71.6*	10.4	53.4	59.2	1.	13.7	36.6	74.3
5547	13	IN 77249RC1-133-3	92.5	C	96.7 B	92.5 C	70.6+	11.2	53.9	54.7	i	15.1	33	73.3
5546	14	IN 78434RB1-10	102.9	A	103.7 A	102.9 A	73.7	10.2	53	55 <b>.</b> 9	1 .	10.8	34	76.3
5549	15	IN 79424M1-20	98.7	B	104.5 A	98.7 B	72.4	10.9	52.4	55.3	ŀ	12.6	33.5	75. 1
5550	16	OH 257	92	C	85.5 D	85.5 D	70.4*	11	55.6 <del>*</del>	47.9	ŀ	15.3	27.5	73.1
5551	17	OH 262	97.2	В	104.5 A	97.2 B	72.0*	10.9	52.4	58.2	1	13.2	35.8	74.7
5552	18	OH 265	97.7	B	107.3 A	97.7 B	72 <b>.</b> i	11	51.7	59.1	ı	13	36.5	74.8
5553	19	OH 285	95	В	92.1 C	92.1 C	71.3*	12.7*	53.8	51.9	1.	14.1	30.8	74
5554	20	DH 286	97.3	В	109.5 A	97.3 B	72.0±	10.4	51.1	61.3	1 -	13.1	38.3.	74.7
5555	21	HART	95	B	98.2 B	95 B	71.3*	11	53.7	54.8	i	14.1	33.1	74
5556	55	SCOTTY	100	A	100 A	100 A	72.8	10.9	52.5	48.0	1	12	27.6	75.5
5557	23	AUBURN	96	В	100.3 A	96 B	71.6*	11 1	53.2	52,3	1	13.6	31.1	74.3
5558	24	TITAN	90	ε	96.9 B	90 C	69.8*	11	54	53.0	1	16.1	31.7	72.6
5559	25	ABE	97.4	В	92.4 C	92.4 C	72.0	12.2*	53.9	51.2	1	13.1	30.2	74.7

Table 2.

1986 CROP

Wheat milling and baking quality data from 1986 Uniform Eastern Soft Red Winter Wheat Nursery samples.

UNIFORM EASTERN RED NURSERY STANDARD = 86302, KNOX 62

9100202

#### WHEAT AND MILLING DATA

LAB	ENTRY	HILLING		COMBINED	TEST		ST.GR.	RED.	FRIABILITY	E. S. I.	MILLABILITY
NO.		QUALITY		QUALITY	WT.	FLOUR		PASSES			
		SCORE	SCORE	SCORE		YIELD	YIELD				•
1.							÷			•	
***	STANDARD - Knox G	2 100 A	100 A	100 A	59.6	33.7	75.2	. 7	27.6	12.1	98. i
- <del>251</del>	BENCHMARK	112 A	110.6A	110.6A	61.6	35.6	76.3	7	28	10.4	114.1
301 1	TRUMBULL	98.4 B	84.1 E	84.1 E	59.8	30 (	76.1	8	27.3	11.9	97.4
305 5	KNOX 62	100 A	100 A	100 A	59.6	33.7	75.2	7	27.6	12.1	98.1
303 3	ORSIS	103.8A	95.2 B	95.2 B	61.1	29, 50		7	29	11.7	104.8
304 4	MD 72004	105.69	102.8A	102.89	60.5	31.10		7	28.8	10.8	107.3
305 2	9021L	96.3 B	95 B	95 B	58. 94		74.4	7	28	12	94.8
306 6	IN 7678862-5-4-7	108.3A	103. 3A	103.3A	60.8	33.4	76.3	7.	29.2	10.6	109.9
307 7	MOW 10501	101.3A	101.4A	101.3A	61	32.7	75.8	7	28.2	11.7	99.3
308 8	IN 77249RC1-133-2	100.7A	100.8A	100.74	60.3	32.3	75.6	7	28	11.9	99.3
309 9	ILL 81-3737	105.5A	94.9 C	94.9 C	60.8	29.20		7	28.2	10.8	107.8
310 10	MOW 11138	99.2 B	103 A	99.2 B	59.1	30.50	75.9	7	29.2	11.6	99
311 11	ND 55-220-76	105.5A	101.5A	101.5A	61.2	33.2	76. 1	7	29	10.5	105.4
312 12	NA SH78-111	96.6 B	95 B	95 B	51.4	30 Q	75.8	7	27 *	11.5	93.2
313 13	1LL 79-1385	102.7A	98.2 B	98.2 B	59.9	29.50	77.3	7	26.4	11.3	104.3
314 14	NA SH76-180	102.7A	99.4 B	99.4 B	59.9	33.1	75.5	7	28.5	11.1	102.3
315 15	PS 840026 .	97.7 B	95.1 B	95. i B	59.9	30, 50	75.3	7	27.8	12.6	95. 9
316 16	COKER 85-42	105.8A	100 A	100 A	60.5	37	75.7	7	29.5	44 7	104.4
317 17	COKER 83-23 (P9323)	112.5A	105. 1A	105. IA	59.7	33.9	77.3	7	30	11.3 9.5	104.4 117.2
318 18	MD 55-217-63	106.8A	99.28	99.2 B	60.1	31.5Q	76.3	7	28.6	10.4	109.3
319 19	KY 83-60	108. 1A	96.6 B	96.6 B	60.5	27.10	77.4	7	29.8	10.4	113.3
320 20	COKER 84-33	101.1A	90.7 €	90.7 C	61	29.30	75.4	7	29.1	11.5	100.7
321 21	COKER 82-28	100.3A	97 8	97 B	61.1	32.1	75.7	7	28.4	11.5	97.9
355 55	OH 257	99 B	93.7 C	93.7 C	59.8	28.80	75.3	7	27.9	11.5	99
323 23	OH 265	104.6A	109.5A	104.6A	60	37.4	76	7	28.9	11.3	102.8
324 24	OH 285	114 A	103.29	103.2A	60.2	29.80	77.4	. 7	29.7	8.7	123.3
325 25	IL 82-3298	102, 8A	94.8 C	94.8 C	61.4	30.60		7	27.5	11.3	102.6
326 26	IL 82-2986	111.8A	94.8 C	94.8 C	61.4	29.20	77 3	7			
327 27	X 1349-10			108.8A	60.6		76.4	7	.28.9	10	117.2
328 28	NA SW76-261			90.9 E		31.50		7	29 97 r	10.5	108.9
329 29	AT 42263-8			104.6A			75. 3	7	27.6	11.1	95.4
330 30	AT 74107-F11-4			101.3A			76.8	7	28.5 28.8	11.8 10.8	105 110. i
331 31	AGC B8 (TRID)	105.4A	105. 2A	105. 2A	59.5	36.5	75 P	7	20.7		
332 32	AGC 89			96.3 B			75.8 75	7	28.1	11.4	105
	- <del>-</del>	*******	-310 D	-U+U D	JJ. J	J7. I	75	7	27.4	12.1	98.6

O1121 O181 E11	airing in	en wokseni	ï
STANDARD =	86302,	KNOX 62	

		STRAIG	HT-GRADE	FLOUR					CAK	E PATENT	FLOUR			
LA NO		r. Ash *	MICI AWRE		TOP GRAIN	1	PROT.	ASH 1	INIT	T FINAL PH	RESPONSI	ELIQUII	CAKE D VOLUME	CAKE SCORE
·				Latte	-				* .		PH/ML/G	LEVEL	<b>X.</b>	
***	10.	9 .39	9 53	17.52										
***	8.9				6 7	1	9.8	.28		1 4.82	2. <del>8</del> 27	130	1080	80
:			- 0484	10100	,	•	7.65	.27	5,6	8 4.84	2.83	130	1048	87
301	11.			17.27*	6	ļ	10.3	.3	5.7	1 4.82	4.571	130	1001 B	a.s.
305	10.			17.52	6	1	9.8	.28		1 4.82	2.827	130	1080	82 80
303	11.			<b>*</b> 17.39	6	1 -	10.3			7 4.81	2.678	130	1079	82
304	10.			17.5	6	1	9.07			5 4.82	2.864	130	1066	84 -
305	10.3	3 .4	53. 3	17.3	5	1	9.44	.27		3 4.81	2.841	120	1060	80
306	10.3	. 39	53.5	17.81	6	1	9.24	.25	5 79	4.84	3.017	120	1671	
307	10.1	. 41	52.8	17.76	7	i	5.13	.3	5. 83		2.618*	150	1071	80
308	10.2	4	53.8	17.34	6	ì	9.11	.29		4.84	2.718	130 120	1041 +	84
309	10.5	.39	. 54.1	17.3	7	1	9.4	.3	5.74		2.838	120	1095 1065	80
310	10.7	.431	53.2	17.58	6	ı	9.43	.3		4.77	2.7	130	1084	80 82
311	10.5	.41	51.5	17.69	7		9.49	26	E 17					
312	11	. 434		17.43	6	,	9.88	.26 .3		4.75	2.973	130	1044 #	84
313	10.7	424		17.48	6	j	9.27	.29	5.74 5.83		2.698	120	1039 *	82
314	9.65		51.2	17.62	6	i i	ŝ. 8	.28			2.782	130	1045	84
315	10.6	-4	53. 1	17.24#	7	İ	9.48	.29		4.83 4.77	2.918 2.745	120 120	1039 * 1065	50 80
316	9.58	.41	56.80	17.45	7						•			•
317	10.7	. 39	51.2	17.71	.7 5	1	8.48	.29		4.84	2.844	120	1110	7 <b>6</b>
318	10.4	.39	51.2	17.71		l .	9.66	.26	5. 78	4.84	2.91	130	1076	62
319	10.9	.41	52.1	17.37	7 6	1	9.32	.28	5.79	4.82	2.592*	120	1049	80
320	10.7	41	49.7	17:08+		1	9.72	.28		4.79	2.645	130	1054	88
	••••	• 14	7301	17.00*	7	1	<b>9.</b> 66	.28	5.87	4. 78	2.59 +	130	1024 *	64
351	10.6	.42#	52. i	17.54	4	1 -	9.86	.31	5.83	4.8	2.774	120	1051	80
322	10.6	.4	53.5	17.43	6	1	9.17	.26	5.88	4.78		120	1029 *	82
323	9.82	.41	52.5	18	7	Í	8. 67	. 27	5.73	4.79		120	1086	80
324 325	10.4	.37	49.3	17.98	6	1	9.58	.27	5.77	4.83		120	1037 *	82
363	9.82	39	53, 3	17.38	7.	1	8. 6	.28	5. 91	4.82		130	1019 +	88
326	11.1	.37	51.4	17.65	,					·				
327	9.42	.39	54.5		6	!	9.96	.27	5. 78		3.005	120	1055 +	80
328	9.88	.43*	52.1	17.84 17.07±	6	1	8.51	.3	5.78			120	1089	88
329	9.82	.37	53	17.65	5 ·	ł	8.95	. 35		4.6		30	1015 *	86
330	10.3	.39	51.8	17.49	6 -	1	8.85 6.85	.26	5.79			20	1080	<b>62</b>
			2110	11.73	6	1	9.25	.28	5.83	4.83	3.124 1	.20	1065	82
31	9.57	. 38	55.4#	17,56	7	i	8.57	.29	5.78	A A7	3. 291 1	20	4405	••
32	10.5	. 38	55. 1÷	17.18*	5	ł	9.43	.27	5.75					84
												LY	1.400	82

1987 CROP UNIFORM EASTERN RED NURSERY

STANDARD = 302, KNOX 62

#### WHEAT AND WILLING DATA

LAB	ENTRY	WILLING	BAKING	COMBINED			ST. GR.			E.\$.I.	WILLABILITY
NO.		QUALITY SCORE	QUALITY SCORE	QUALITY SCORE	WT.	FLOUR YIELD	FLOUR YIELD	PASSES	•		
		***************************************	440.11			,					
111	STANDARD - Knox 62	100 A	100 A	100 A	62	31.5	75.9	8	27	12	107.9
***	BENCHMARK	101.3A,	114.2A	101.34	61.6	35.8	76.3	7	28	10.4	107.9
301 A	TRUMBULL	97.9 B	50.δ F	50.6 F	60.6Q	32.8	75.8	7 -	21	11.4	104.9
302 B	KNOX 62	100 A	100 A	100 A	. 82	31.5	75.9	8	27	12	107.9
303 C	OASIS	96.7 8	18.9 F	78.9 F	62.6	29 Q	76.1	. 1	27.9	11.6	103.2
304 BFA	TWAIN (NA SW78-111)	91.3 C	84.6 E	84.6 E	62.4	31.4	75.5	8	26.19	12.4	92.7 *
305 8FC	COKER 9733 (84-33)	97.9 8	84.3 E	84.3 E	62.5	29.5Q	75.4	1	28.5	11.3	104.9
306 IB	DYNASTY (OH 265)	101.1A	81.5 E	81.5 E	61.6	36.3	75.8	1	28.2	11.3	107.2
307 FR	PIONEER 2555 (XW 546)	102.6A	98.2 B	98.2 B	66.60		76.2	1	28.6	10.3	110.5
308 BFH	PACER (HY. EXP. LE-10)	96.5 B	112.9A	96.6 B	_	32.9	75.4	1	28.1	11.3	102.1
308 BEU	LINCLON (NA-76-180)	100 A	86.1 D	86.1 D	60.50		75.5	1	27.9	11	107.9
310 EFI	AR 74107-F11-4	101.7A	73.1 F	73.1 F	61.18	29.30	76.7	7	28	10.7	113.1
311 BGS	IL 83-3298	97. 8 :	72.8 F	72.8 F	62.6	31	75.7	8	26.6*	12.1	102.4
312 EF	WI X 1336-8	100.5A	115 A	100.5A	60 Q	35.2	76.7	7	28.8	9.9	108.5
313 BDX	IL 81-3737	95.4 B	69.9 F	69.9 F	61.6	27.20	76.4	8	27.1	11.2	103.2
314 BGY	KA-83-00	101 A	. 78 F	78 F		28.2Q	71.2	1 .	28.8	9.7	112.5
315 SGT	IL 82-2986	102.1A	87.9 D	87.9 D	52.1	29.10	77.1	1	28	10	112.8
316 IC	OH 285	107.8A	99 B	99 8		29.3Q	77.4	7	29.3	9	123.2
317 BFJ	AR 42263-8	444 44	81.9 E	81.9 E	60.89	•	75.1	1	27.5	11.9	108.4
318 BF0	AGC-107	101.5A	83.8 E	83.8 E	61 \$	36.6	76.4	8	27.7	11.2	108.3
319 BFP	AGC-108	94.1 0	77 F	17 F		34.3	75.1	8	26.4*	12.4	97.3 \$
320 IE	OH 262	93.1 C	102.7A	93.1 C	59.79	38.6	75 \$	8	27.3	12.4	94.1 #
321 IO	OH 286	97.1 B	89.4 D	89.4 D	59.50		75.5	1	27.8	11.5	101.2
322 BGU	IN. 771454-C1-2-9-69	93.5 C	65.1 F	65.1 F	61.18		15.4	8	27.1	12	97.7 *
323 OT	IN. 79404-61-28-2	99.9 8	88.8 D	88.8 0	62.4	31.8	75.9	1		10.5	107.1
324 807	IN. 77249-RC1-133-3	91.8 C	85.1 D	85.1 D	60.91		75.3	8	26.5*		.~ 93.8 *
325 86N	MD 75191-80	103.6A	69.1 F	69.1 F	60.30	32.1	76.5	1	28.4	9.9	115.5
326 8GJ	MD 73019-26	89.4 D		84.6 E	59.76		74.5*	8		12.9	90.5 *
327 BGK	ND 73025-51	102.6A	87.6 D		59 Q		17.5	8		9.8	113.3
328 BGL	ND 73065-03	87.7 D	92.3 C		59 Q		74.9*	8		12.1	87.4 Q
329 BFV	COKER 86-30	100.1A	90.3 C		61.9		76.2	i		10.9	104.2
330 B6W	PS 640061	104.8A	85.1 D	85.1 D	60.7Q	33	11.1	7	29.1	9.8	116.5
331 BFX		97 8			81.1*		78	i		11.4	99.8
332 BFY	COKER 86-38		93.9 €			27.10		7		10.2	117.7
333 BGV	PS 840024	101.2A	57.3 F			•	76.6	i		10.2	112.4
	** *****	IVIIIA	J (1) (	9713 F	u J . 4	T0.34	10.0	· ' .	20.3	10	114.4

1987 CROP

### UNIFORM EASTERN RED NURSERY STANDARD = 302, KNOX 62

#### STRAIGHT-GRADE FLOUR

LAB	FLOUR	FLOU	R	MICRO	COOKIE	TOP
NO.	PROTEIN	ASH		AVRC	DIANETER	GRAIN
	\$	3		1	CN.	winan
1.1		-		•	•	
111	10	.34		49.8	17.84	7
***	8.899	.35		51.3	18.35	1
201		3.0			40.0.0	•
301 302	11.48Q 10	.36		51.1	16.8 <b>Q</b> 17.84	3 7
303	18.22	.34		49.8 50.8	17.39*	6
304	9.84	,4	Q	50.5	17.49*	6
305	9.85	.38		48.7	17.4 *	
305	9.43	.37	•	51.2	17.42*	. 5
	0110			VIII		. •
307	8.81	.38	Q	50.7	17.74	6
308	8.95	.39		49.7	18.1	5
309	8.94	.36	ŧ	49.7	17.45*	6
310	9.67	.36	ŧ	50.5	17.2 Q	4
311	9.16	.36	1	52.6 *	17.260	i,
312	9.01	. 4 1	Ŷ	49.5	18.14	6
242	A 61	20			. 43 450	
313	9.64 9.4	.39	Q	50.7	17.130	3
314 315	9.84	. 4	Q	50.5	17.3 Q	3
316	9.8	.38	Q	51.2 49.1	17.6 17.77	4 8
317	8.58	.33	٠	51.9 *	17.45*	5
318	8.85	.37	1	52.9 *	17.54*	5
710	4:42	131	·	32.3	111347	J
319	9.85	.37		51.9 *	17.37*	À
320	8.72	. 4	Q.	50.1	17.87	4
321	9.1	.39	Q	49.7	17.53*	3
322	10.2	.39	Q	50.5	17.040	10
323	9.15	.38	Q	49.8	17.52*	ă.
324	9.37	.39	ę.	51.8 *	17.53*	<b>4</b>
325	8.38	.36		52.1 *	45 450	
326	8.25	.39		51.6 *	17.15Q 17.5 *	5 5
327	8.66	. 4	Q	49.1	17.46*	4
328	9.05	.42	Q	51	17.46-	2
329	8.58	.41	Q	53.8 Q	17.74	8
330	8.98	.39	è	50	17.44*	4
		.,,	٦	**	11447	7
331	7.94	.42	Q	53.1 Q	17.61	4
332	9.8	.36	ì	49.4	17.66	3
333	9.31	.39	Q	51.1	16.829	4
334	9.39	.42	Q	51.6 *	17.6	4

988 CROP

ILFORM EASTERN RED NURSERY TANDARD = 302, KNOX 62

## WHEAT AND MILLING DATA

LAB NO.	ENTRY	MILLING QUALITY SCORE	BAKING QUALITY SCORE	CONBINED QUALITY SCORE		BREAK FLOUR YIELD	ST. GŔ. FLOUR YIELD	RED Passes	FRIABILITY	E.S.I.	YTL1881LITY	
				400 1	C 1 0	27.8	76.0	8	28.3	11.7	98.4	
***	STANDARD - Knox 62	100 A	100 A	100 A		35.8	16.3	1	28	10.4	98.4	
***	BENCHMARK	103.1A	103.2A	103.1A	61.6		76.0	8	28.3	11.7	98.4	
302 2	KNOX 62	100 A	100 A	100 A	61.8	27.8	15.7	8	26.3	12.0	95	
	OASIS	97 B	89.20	89.2 D	61.8	25.8Q	16.0	7	26.7	11.3	102.3	
303 3	IL 82-3298	102.4A	81.30	87.3 0	61.8	27.5	15.1	<del></del>	27.5	11.5	104.4	
304 4	COKER P9733	103.8A	84.3 E	84.3 E	81.8	27.7	19.1	,	21,14	•		
305 5	CONTRACTOR	-					70 1	8	26.5	11	103.4	
	AGC-107	104.4A	98 8	96 B			76.1	1	21.7	10.4	104	,
306 6	COKER 86-35	105.2A	94.9 C	94.9 C	60.9*		16.5	1	27.0	11.4	105	
307 7	COKER 86-38	102.6A	88.1 Û	88.1 D	61.8	24.0Q	78.1	,	27.3	P 11.4	103.2	
308 8		104.3A	100.2A	100.2A	60.29		75.8	,	28.8	10.0	113.4	
309 9	OH 286	111.7A	98.4 8	98.4 8	60.69	34.3	16.8	1	70.0	,,,,,		
310 10	PIONEER 2555	••••							27.7	10.8	110.9	
		108.84	80.5 E	80.5 E	61.9	29.4	76.7	1		10.7	109.3	
311 11	HANCOCK	196.1A	89.8 0	89.8 D	80.9*	26.9*	76.5	7	27.3		94.2	
312 12	NA 5884-345	97 8	87 0	87 D	61 *	28.4	75.4	7	26.9	11.9	95.9	
313 13	PACER	98.6 8	87 D	87 D	61.5	28.9	75.4	1	26.5	12.2	98.5	
314 14	77145401-2-9-6914		94.8 C	94.8 C	61.5	28.5	15.7	1	26.9	11.2	20.3	
315 15	7940461-26-2	100.2A	. 3410 0									
21	•		75.7 F	75.7 F	59.30	30.0	18.8	1	27.8	9.73	112.2	
316 16	ND 75191-80	108.2A		88.2 0	59.80		76.8	1	27.8	10.2	107.6	
317 17	MD 73025-51	105.7A	88.2 0	84.2 E	60.40		74.6*	8	25.7*	12.1	86.2 *	• *
318 18	ND 73065-03	92.5 C	84.2 E	68.6 F	61.6	29.0	74.20	8	25.8*	13.5*	79.6 Q	
319 19	AR 26415	88.2 D	68.6 F		61.1*		15.2	8	28.1	12.3	91.9	
320 20	IL 84-2518	96.1 8	79.4 F	79.4 F	VIII'							
				65 6	61.9	24.70	76.9	7	27.5	10.6	104.3	
321 21	IL 82-2766	102.6A	93 C	93 C	59.8Q		76.1	7	27	11,3	103.6	
322 22	AL 840169	103.6A	85.6 D	85.60			75.3	1	26.9	12.0	93.5	
323 23	AL 850046	97.7 B	87.4 D	87.4 D	60.8		74.6*	1	26.0	12.6	90.8	
324 24	2 U 51	95.1 8	70.1 F	70.1 F	61.4	28.5		. 8	25.6Q	12.1	92.1	
	2 C 21	94,1 C	· 69.8 F	69.8 F	61 Z	24.3	<b>á</b> 1313	•		, •		
325 25	2 0 21	•				1	78	i	28.5	9.38	108.8	
	WI X1656-1	104.2A	91 C	91 C		26.4		i	28.6	9.8	110.5	
326 26	COKER 86-22	107.8A	111.84	107.8A		29.3		i	27.4	11.9	96.8	
327 27		99.3 B	86.9 D	86.9 D	62	28.7	75.8		27.0	11.6	95.7	
328 28	NO 9965	97.8 B	74.3 F	74.3 F		28.3		8	27.1	11.6	100.4	•
329 29	OH 331	99.8 B	71.6 F	71.8 F	60.7	25.8	Q 75.8	1	21.1	,,,,,		
330 30	OH 374	JU10 H						_	44.4	9.94	110.8	
		108.1A	91.1 C	91.1 C	60.5	Q 30.2		1	28.0		105.2	
331 31	OH 394		93.2 C	93.2 C		31.0		1	27.6	10.8	112.7	
332 32	1942441-20-2-74	104.8A	95.9 B		61.6					10.1	101.8	
333 33	KA 83-38	111.34	88.2 D		60.8			7	27.5	11.3	101.0	•
334 34	NO10501	102.8A	90.2 8					-9			14	L

Table 5. Wheat milling and baking quality data from 1989 Illinois Advanced Nursery samples.

DR. F.L. KOLB URBANNA, ILLINOIS

STD = #2005, SCOTTY

LA8 NO		ENTRY		MILLING QUALITY SCORE	QUALITY	COMBINED QUALITY SCORE	NICRO T.W. KG/HL	EQUIV.	FLOUR YIELD		MICRO AWRC	COOKIE DIAM.		
****		TANDARD ~ ENCHMARK	Seatty			100.0 A 97.4 8	79.6 78.6	46.8 61.4	78.6 75.1 *	9.87 1.87	54.3 52.9	18.14 18.21	3	
2003	1 0	ALDWELL		101.2 A	88.8 D	88.80	79.2	53.6	77.0	B 07	67.0	17.83 *		
2004	2 A	JBURN '		99.7 8		79.4 F	80.7	48.2	76.0	9.76		17.08 Q		
2005	3 \$	COTTY			100.0 A		79.6		76.6	9.87		18.14	3	
2006	4 88	CKER		96.9 B	101.4 A	98.98	75.8	57.0	76.4	1.67		18.31		
2007	5 C	ARDINAL		103.5 A	93.0 €			51.8	11.1	8 44			3	
										• • • • • • • • • • • • • • • • • • • •				
2008	1000	NASTY		98.5 B	91.6 C	91.5 C	79.0	55.5	75,4 *	9.37	56.7 *	17.9	2	
2009	1 1	the state of the s		98.4 8	82.8 E	82.8 E	78.2		75.5			17.68 *		
2010		LAND		103.2 A	99.3 8	99.3 B	79.1	51.4	76.6	8.69	55.2		4	
2011		ARX		91.3 C	67.5 F	87.5 F	78.6	45.8	74.7 *	9.70	58 Q	17.25 Q	3	
2012	10 PI	0 2555		107.9 A	98.6 B	98.88	78.8	58.2	78.1	9.18	56.1 *		4	
0040	44.6*	19 P S												
2013		0 2548				65.7 F		50.8	74.8 *	9.69	58.7 Q	16.93 Q	. 2	
2014		R 133-3		95.2 8	92.8 C			50.2	74.5 *	8.92	56.5 *	17.93	3.	
2015	13.90			104.8 A	and the second second	96.9.0		50.3	76.7	9.42	56.6 *	17.66 *	3 -	
2016		82-3298 L 84-2518-1		102.0 A	67.9 F	67.9 F		46.3	76.9	9.21		17.22 Q		
2011	19 16	L. 04-2310-1		92.3 C	77.4 E	77.4 F	78.2	51.5	73.8 Q	9.92	57.7 Q	17.4 Q	2	
2018	16 111	84-3010		100 ( )	10 0 5	10.0.								
		L 84-3511		100.1 A		79.2 F			76.1	8.52		17,43 Q		
2020		84-4046		101.1 A	81.3 E	81.3 E		52.6	76.1	8.95		17.52 Q		
2021		84-2191-1	$N_{X,X} \cdot (x, x, y) = xx_{X,X} \cdot (x, y) + \dots = x_{X,X} \cdot (x, y) \cdot (x, y)$	98.3 8					75.1 *		at a second	17.77 *		
2022		84-3659-1		99.6 8		94.7 C		54.2		8.72	57.7 Q		3	
2422		. 04 3933 (		33.0 0	89.5 F	89.5 F	78.4	50.8	75.8	8.75	59.1 Q	17.14 Q	3	
2023	21 111	84-5464	100	100.5 A	87.10	81 1 0	70 6	EA: A	30.0	0 00			_	
2024		85-2872	1	96.3 8	62.0 F	87.10			76.0	8.89		17.81 *		
2025		85-3132-1			76.3 F				74.9 *			16.65 Q		
2026		85-5865		97.3 B	78.3 F				74.1 Q			17.33 Q		
2027	25 CAL			105.5 A	86.1 D	86.10			75.2 * ·			17.36 Q		•
					00.1 8	00.10	10.4	34.5	77.4	9.21	38.2 W	17.83 *	3	
2028	26 CAR	DINAL		106.1 A	93.3 1	93.8 0	79.0 5	32.2	11.4	9 62	CC 0 ¥	17.89 #	3	
2029	27 KNG			94.9 C	66.2 F	66.2 F						17.15 Q		
2030		82-3298		103.0 A	81.0 E	81.0 E						17.15 Q 17.87 *		
2031		84-3565-1		99.6 B						8.29		17.94		
2032		77-3457		99.08	15.6 F							17.36 Q		
					***			7,3		v.34	ים ייט על	i1.•30 A	I.	

#### Exhibit E

# Statement of the Basis of Applicant's Ownership

The complex cross, generation advance, selection, testing and evaluation, repurification, and multiplication were all performed by the applicant breeder (F. L. Kolb) or his predecessor (C. M. Brown) or technical assistants on the property of the Illinois Agricultural Experiment Station. The variety is intended for use as a public variety in the United States.

The employee's rights, if any, are vested in the University as a condition of employment.